

Injective envelopes and flat covers of Matlis reflexive modules

Richard G. Belshoff

Department of Mathematics, Southwest Missouri State University, Springfield, MO 65804, USA

Jinzhong Xu*

Department of Mathematics, Suzhou University, Suzhou 215006, People's Republic of China

Communicated by M. Barr

Received 20 January 1991

Revised 13 August 1991

Abstract

Belshoff, R.G. and J. Xu. Injective envelopes and flat covers of Matlis reflexive modules, *Journal of Pure and Applied Algebra* 79 (1992) 205–215.

We show that for a commutative noetherian local ring R , every Matlis reflexive R -module has a reflexive injective envelope if and only if every Matlis reflexive R -module has a reflexive flat cover. This occurs if and only if R is complete and has Krull dimension less than or equal to 1. We also exhibit a family of Matlis reflexive R -modules whose injective envelopes are not Matlis reflexive.

1. Introduction

In this paper R will denote a commutative noetherian ring with identity. If M is an R -module, then $E_R(M)$ or just $E(M)$ will denote an injective envelope of M as an R -module.

The notion of a flat cover of a module was introduced by Enochs in [5]. A flat cover of an R -module M consists of a flat module F and a linear map $\sigma : F \rightarrow M$ with the following two properties:

Correspondence to: R.G. Belshoff, Department of Mathematics, Southwest Missouri State University, Springfield, MO 65804-0094, USA. Email: rgb865f@smsvma.bitnet.

* Current address: Department of Mathematics, University of Kentucky, Lexington, KY 40506-0027, USA.

(i) For any flat module F' and any linear map $\sigma' : F' \rightarrow M$ there is a linear map $g : F' \rightarrow F$ such that

$$\begin{array}{ccc} & F' & \\ g \swarrow & \downarrow \sigma' & \\ F & \xrightarrow{\sigma} & M \end{array}$$

commutes.

(ii) Any linear $g : F \rightarrow F$ such that

$$\begin{array}{ccc} & F & \\ g \swarrow & \downarrow \sigma & \\ F & \xrightarrow{\sigma} & M \end{array}$$

commutes is an automorphism.

If $\sigma : F \rightarrow M$ satisfies (i) it is called a flat precover. Clearly such a σ is always reflexive. In [6, Proposition 1.2, p. 180] it was shown that if R is a complete local ring, then every Matlis reflexive R -module has a flat cover.

For a local ring (R, \mathfrak{m}) with residue field $k = R/\mathfrak{m}$, let $E = E_R(k)$. Then M'' denotes the Matlis dual $\text{Hom}_R(M, E)$ of the R -module M , and M is said to be Matlis reflexive if the canonical map $\tau_M : M \rightarrow M''$ is an isomorphism. It is easy to check directly that τ_M is always injective. In this paper ‘reflexive’ will always mean Matlis reflexive. We need the following results of Matlis.

(a) Let \hat{R} be the \mathfrak{m} -adic completion of R . Then by [9, Theorem 3.7, p. 522],

$$\text{Hom}_R(E, E) \cong \hat{R}.$$

(b) Let R be a complete local ring. Then finitely generated and artinian R -modules are reflexive. Also, M is an artinian (noetherian) R -module if and only if M'' is noetherian (artinian) [9, Corollary 4.3, p. 528]. In particular, $R'' \cong E$ is artinian, $E'' \cong R$ and both are reflexive.

The main results of Matlis [9] are nicely summarized in Matsumara’s book [11]. See also the book by Sharpe and Vámos [13].

We note for use later the following general ‘change of ring’ facts. Let R be any ring and let $I \subset R$ be an ideal. Given an R/I -module M , we can regard M canonically as an R -module. Suppose now that E is an injective R -module. Let

$$\text{ann}_E(I) = \{x \in E \mid Ix = 0\}.$$

Then $\text{ann}_E(I)$ is a submodule of E , and it is clear that $\text{ann}_E(I)$ is also an R/I -module. It is not hard to see that $\text{ann}_E(I)$ is an injective R/I -module.

Lemma 1.1. *Suppose M is an R/I -module and $E = E_R(M)$. Then*

$$E_{R/I}(M) = \text{ann}_E(I).$$

Proof. Since M is an R/I -module, $M \subset \text{ann}_E(I)$. We have just noted that $\text{ann}_E(I)$ is an injective R/I -module. It remains to be shown that $\text{ann}_E(I)$ is an essential extension of M as an R/I -module. But this is true since as R -modules it is clear that $\text{ann}_E(I)$ is an essential extension of M . \square

In particular, for a local ring (R, \mathfrak{m}) with residue field $k = R/\mathfrak{m}$, if $E = E_R(k)$, then $E_{R/I}(k) = \text{ann}_E(I)$. But if M is an R/I -module and is regarded canonically as an R -module, then

$$\text{Hom}_R(M, E) = \text{Hom}_{R/I}(M, E_{R/I}(k)),$$

and so M^ν is unambiguous. This means that we can think of M as an R -module or as an R/I -module and M^ν means the same thing. Also note that $IM^\nu = 0$, so $M^{\nu\nu}$ is unambiguous. This means then that such a module M is reflexive as an R -module if and only if it is reflexive as an R/I -module.

2. The injective envelope of a Matlis reflexive module

Let (R, \mathfrak{m}) be a local ring. If M is an R -module and $S \subset M$ a submodule, consider the commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & S & \longrightarrow & M & \longrightarrow & (M/S) \longrightarrow 0 \\ & & \downarrow \tau_S & & \downarrow \tau_M & & \downarrow \tau_{(M/S)} \\ 0 & \longrightarrow & S^{\nu\nu} & \longrightarrow & M^{\nu\nu} & \longrightarrow & (M/S)^{\nu\nu} \longrightarrow 0 \end{array}$$

with injective vertical maps. A diagram chase using the snake lemma shows that M is reflexive if and only if S and M/S are reflexive. A finite direct sum $\bigoplus M_i$ is reflexive if and only if each M_i is reflexive. In particular, if $E(M)$ is reflexive, then M , being a submodule, is also reflexive. But we note that the converse is not true in general. For example, if k is a field then $R = k[[x, y]]$ is a complete local ring and so is Matlis reflexive. We will show that $E(R)$, the field of fractions of R , is not a Matlis reflexive R -module.

Given a reflexive module M , in this section we establish conditions under which the injective envelope $E(M)$ is also reflexive module.

Proposition 2.1. *If (R, \mathfrak{m}) is a complete local ring and M is an artinian R -module, then $E(M)$ is reflexive.*

Proof. We have already noted that if $k = R/\mathfrak{m}$, then $E(k)$ is reflexive. Since $E(M) \cong E(k)^n$ for some n [9, Proof of Corollary 4.3], it follows that $E(M)$ is reflexive. \square

Lemma 2.2. *Let R be any local ring, M an R -module, and $S \subset M$ a submodule. If $E(S)$ and $E(M/S)$ are both reflexive, then $E(M)$ is reflexive.*

Proof. There is an obvious injection $M \rightarrow E(S) \oplus E(M/S)$ which makes the diagram

$$\begin{array}{ccccccc}
 0 & \longrightarrow & S & \longrightarrow & M & \longrightarrow & M/S \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & E(S) & \longrightarrow & E(S) \oplus E(M/S) & \longrightarrow & E(M/S) \longrightarrow 0
 \end{array}$$

commute. It follows that there is an injection $E(M) \rightarrow E(S) \oplus E(M/S)$. Thus $E(M)$ is a direct summand of a reflexive module and therefore $E(M)$ is reflexive. \square

We begin by assuming that R is a domain.

Proposition 2.3. *Suppose R is a complete local domain and $\dim R = 1$. Then M is a reflexive R -module if and only if $E(M)$ is reflexive.*

Proof. We only need to show that if M is reflexive, then $E(M)$ is reflexive. For an R -module M , let tM denote the torsion submodule of M . Replacing S in Lemma 2.2 by tM , we get that $E(M)$ is a direct summand of $E(tM) \oplus E(M/tM)$. Since M is reflexive, the submodule tM is as well, hence so is M/tM . So it is enough to show that $E(M)$ is reflexive in both cases— M torsion and M torsion-free. We do this using the following lemmas.

Lemma 2.4. *Let (R, \mathfrak{m}) be a complete local domain with $\dim R = 1$. If S is a finitely generated torsion-free R -module, then $E(S)$ is reflexive.*

Proof. We will denote the field of fractions of R by Q and will let $k = R/\mathfrak{m}$ be the residue field. Then S embeds in a finitely generated free R -module, say R^n for some n [12, Lemma 4.31]. Noting that $Q = E(R)$, we have $E(S) \subset Q^n$, so it is enough to show that Q is reflexive. And since R is a finitely generated R -module, to show that Q is reflexive it is enough to show that Q/R is artinian [9, Corollary 4.3 and Snake Lemma]. But this follows from [10, Theorem 1]. \square

Lemma 2.5. *Let R be a complete local domain with $\dim R = 1$. If M is a torsion-free R -module and is reflexive, then $E(M)$ is reflexive.*

Proof. Since M is reflexive, by [6, Proposition 1.3, p. 181] there exists a finitely generated submodule $S \subset M$ with M/S artinian. But $E(M/S)$ is reflexive by Proposition 2.1, and $E(S)$ is reflexive by Lemma 2.4. Therefore, $E(M)$ is reflexive by Lemma 2.2. \square

Lemma 2.6. *Let R be a complete local domain with $\dim R = 1$. If S is a finitely generated torsion R -module, then S is artinian and so $E(S)$ is reflexive.*

Proof. We first consider the case where $S = Rx$ is cyclic. Then $\text{ann}(x) \neq 0$ since S is torsion, hence $\dim(R/\text{ann}(x)) = 0$, and so $R/\text{ann}(x) \cong S$ is artinian. Therefore, $E(S)$ is reflexive by Proposition 2.1.

Next suppose that $S = Rx_1 + \cdots + Rx_r$, and let $T = Rx_1 \oplus \cdots \oplus Rx_r$. There is an obvious surjection $\phi : T \rightarrow S$, but T is artinian by the above argument, and therefore S is artinian. \square

Lemma 2.7. *Let R be a complete local domain with $\dim R = 1$. If M is a torsion R -module and is reflexive, then $E(M)$ is reflexive.*

Proof. Since M is reflexive, we have a finitely generated submodule $S \subset M$ with M/S artinian. Lemma 2.6 and Proposition 2.1 show that $E(S)$ and $E(M/S)$ are reflexive, and so $E(M)$ is reflexive by Lemma 2.2. \square

This completes the proof of Proposition 2.3. \square

Proposition 2.8. *Suppose R is a complete local domain with field of fractions Q . Then the following statements are equivalent:*

- (1) Q is reflexive.
- (2) Q/R is artinian.
- (3) $\dim R \leq 1$.

Proof. (1) \Leftrightarrow (2) If Q/R is artinian, then Q has a finitely generated submodule—namely R —with an artinian quotient, and so Q is reflexive.

Conversely, if Q/R is not artinian then let $S \subset Q$ be any finitely generated submodule. To show that Q is not reflexive it suffices by [6, Proposition 1.3] to show that Q/S is not artinian.

Suppose that S is generated by s_1, \dots, s_n . Then for each i we have $s_i = f_i/g_i$, where $f_i, g_i \in R$, and $g_i \neq 0$. Setting $y = 1/(g_1 \cdots g_n)$, we have that $S \subset Ry$. But the map $R \rightarrow Ry$ sending 1 to $1y$ is bijective, and so $R \cong Ry$. Hence there is a well-defined epimorphism $\phi : Q/S \rightarrow Q/R$. So Q/S is not artinian.

(2) \Leftrightarrow (3) This follows from [10, Theorem 1, p. 571]. \square

Remark. The proposition supplies plenty of examples of reflexive modules whose

injective envelopes are not reflexive—any complete local domain of dimension at least two (regarded as a module over itself) fits the bill.

Now we drop the assumption that R is a domain.

Theorem 2.9. *Suppose (R, \mathfrak{m}) is a complete local ring with $\dim R \leq 1$. Then an R -module M is reflexive if and only if $E(M)$ is reflexive.*

Proof. First assume $\dim R = 0$. If M is reflexive, then there is a finitely generated submodule $S \subset M$ such that M/S is artinian. But then the finitely generated module S is also an artinian module since R is an artinian ring. So both $E(S)$ and $E(M/S)$ are reflexive by Proposition 2.1, and therefore $E(M)$ is reflexive.

Now assume $\dim R = 1$. Let M be a reflexive R -module. Then for any finitely generated R -module S we can write

$$E(S) = \bigoplus_{\mathfrak{p} \in \operatorname{Spec}(R)} \mu_0(\mathfrak{p}, S) E(R/\mathfrak{p}), \quad (1)$$

where $\mu_0(\mathfrak{p}, S) < \infty$ for all primes \mathfrak{p} [3, Lemma 2.7, p. 11]. Since the prime ideals of R are the maximal ideal \mathfrak{m} and a finite number of minimal prime ideals $\mathfrak{p}_1, \dots, \mathfrak{p}_r$, (1) is a finite direct sum. So to prove that $E(S)$ is reflexive it is enough to prove that $E(R/\mathfrak{p})$ is reflexive whenever \mathfrak{p} is a minimal prime ideal of R . But if \mathfrak{p} is minimal, then $\dim R_{\mathfrak{p}} = 0$, so in this case $R_{\mathfrak{p}}$ is an artinian local ring with residue field $k(\mathfrak{p}) = R_{\mathfrak{p}}/\mathfrak{p}_{\mathfrak{p}}$. From [9] (or [11, Theorem 18.4(vi)]) we know that

$$E_R(R/\mathfrak{p}) = E_{R_{\mathfrak{p}}}(k(\mathfrak{p})).$$

Since the Matlis dual of an artinian is finitely generated [9, Corollary 4.3], we have that $R_{\mathfrak{p}}^{\vee} \cong E_{R_{\mathfrak{p}}}(k(\mathfrak{p})) \cong E_R(R/\mathfrak{p})$ is a finitely generated $R_{\mathfrak{p}}$ -module. And since $\dim R_{\mathfrak{p}} = 0$, $E_R(R/\mathfrak{p})$ has finite length over $R_{\mathfrak{p}}$. This means that there is a composition series for $E_R(R/\mathfrak{p})$,

$$0 \subset A_0 \subset \cdots \subset A_n = E_R(R/\mathfrak{p}), \quad (2)$$

where each factor A_i/A_{i-1} is a simple $R_{\mathfrak{p}}$ -module. So each factor must be $k(\mathfrak{p})$. Recall that $k(\mathfrak{p})$ is naturally isomorphic to Q , the field of fractions of the domain R/\mathfrak{p} . Since $Q = E_{R/\mathfrak{p}}(R/\mathfrak{p})$ is a reflexive R/\mathfrak{p} -module by the domain case (Proposition 2.8), Q is also reflexive as an R -module. Because each factor in the composition series (2) is reflexive as an R -module, $E_R(R/\mathfrak{p})$ is reflexive as an R -module.

This shows that $E(S)$ is reflexive if $S \subset M$ is any finitely generated submodule of M with M/S artinian. But $E(M/S)$ is also reflexive by Proposition 2.1. Hence by Lemma 2.2, $E(M)$ is reflexive. \square

3. The flat cover of a Matlis reflexive module

Let (R, \mathfrak{m}) be a complete local ring. We have already noted that every Matlis reflexive R -module has a flat cover [6, Proposition 1.2]. We now ask whether the flat cover of a reflexive module is reflexive. Because R is a noetherian local ring, it is semi-perfect and every finitely generated R -module has a projective cover ([2, Theorem 2.1] or [12, pp. 135–136]).

Proposition 3.1. *Let $\phi : P \rightarrow M \rightarrow 0$ be the projective cover of the finitely generated R -module M . Then (ϕ, P) is the flat cover of M .*

Proof. (i) For any flat R -module F and homomorphism $f : F \rightarrow M$, we have the diagram

$$\begin{array}{ccccc} & & F^\nu & & \\ & \nearrow g & \uparrow f^* & & \\ P^\nu & \xleftarrow{\phi^*} & M^\nu & \xleftarrow{\quad} & 0, \end{array}$$

where ϕ^* is an injection, and (by [7] or [8, Exercise 8(b), p. 614]) F^ν is an injective module. Hence there is a homomorphism $g : P^\nu \rightarrow F^\nu$ such that $g\phi^* = f^*$. Note that P is finitely generated and P, M are reflexive. We have the diagram

$$\begin{array}{ccccc} & & F^{\nu\nu} & & \\ & \nearrow g^* & \downarrow f^{**} & \nwarrow \tau_F & \\ & P^{\nu\nu} & \xrightarrow{\phi^{**}} & M^{\nu\nu} & \xleftarrow{f} F \\ & \nwarrow \tau_P & \nearrow g_1 & \nwarrow \tau_M & \\ & P & \xrightarrow{\phi} & M & \end{array}$$

We know that τ_P and τ_M are isomorphisms. Set $g_1 = \tau_P^{-1} g^* \tau_F$. A diagram chase shows that $f = \phi g_1$. Therefore, $\phi : P \rightarrow M \rightarrow 0$ is a flat precover of M .

(ii) For any endomorphism $f : P \rightarrow P$ such that $\phi = \phi f$, we need to prove that f is an automorphism of P . But $\ker(\phi)$ is a superfluous submodule of P , and so f is epic. Hence f is an automorphism because P is a noetherian module. \square

We will need the following result due to Ishikawa.

Lemma 3.2. *If E and E' are injective R -modules, then $\text{Hom}_R(E, E')$ is a flat R -module.*

Proof. See [7, Theorem 1.5, p. 293]. \square

For the general case, we have the following result.

Proposition 3.3. *Let M be a reflexive, $F \xrightarrow{\phi} M$ be the flat cover of M , and let $\tau_F : F \rightarrow F^{\nu\nu}$ be the canonical injection. Then F is a direct summand of $F^{\nu\nu}$.*

Proof. Since F is the flat cover of M , ϕ is epic. We have the following diagrams:

$$\begin{array}{ccc} F & \xrightarrow{\phi} & M \\ \tau_F \downarrow & & \downarrow \tau_M \\ F^{\nu\nu} & \xrightarrow{\phi^{**}} & M^{\nu\nu} \end{array} \quad \begin{array}{ccc} F & \xrightarrow{\tau_F} & F^{\nu\nu} \\ & \searrow \beta & \downarrow \tau_M^{-1} \phi^{**} \\ & & M \longrightarrow 0 \end{array}$$

Since $F^{\nu\nu}$ is flat (Lemma 3.2) and $\phi : F \rightarrow M$ is the flat cover, there exists a homomorphism $\beta : F^{\nu\nu} \rightarrow F$ such that $\tau_M^{-1} \phi^{**} = \phi \beta$, and then $\phi \beta \tau_F = \tau_M^{-1} \phi^{**} \tau_F = \phi$. Therefore, $\beta \tau_F : F \rightarrow F$ is an automorphism of F and F is isomorphic to a direct summand of $F^{\nu\nu}$. \square

4. The main result

Theorem 4.1. *Let (R, \mathfrak{m}) be a noetherian local ring. Then the following statements are equivalent:*

- (1) *Every reflexive R -module has a reflexive injective envelope.*
- (2) *Every reflexive R -module has a reflexive flat cover.*
- (3) *R is complete and $\dim R \leq 1$.*

Furthermore, suppose these conditions hold and M is a reflexive R -module. Let E be injective and F flat. Then $M \rightarrow E$ is an injective envelope of M if and only if $E^\nu \rightarrow M^\nu$ is a flat cover of M^ν . Similarly, $F \rightarrow M$ is a flat cover of M if and only if $M^\nu \rightarrow F^\nu$ is an injective envelope of M^ν .

Proof. (1) \Rightarrow (2) Let M be reflexive. Then M^ν and $E := E(M^\nu)$ are reflexive. Let $0 \rightarrow M^\nu \xrightarrow{h} E$ be the injective envelope of M^ν . Let F be any flat R -module and let $f : F \rightarrow M^{\nu\nu}$ be an R -module homomorphism. Then we have the following diagram

$$\begin{array}{ccc} & & F \\ & \searrow g_1 & \downarrow f \\ E^\nu & \xrightarrow{\phi} & M^{\nu\nu} \longrightarrow 0 \end{array}$$

where $\phi = h^*$, and there exists a linear map g_1 such that $\phi g_1 = f$. Since E^ν is flat by Lemma 3.2, this shows that (E^ν, ϕ) is a flat precover of $M^{\nu\nu}$.

For any linear map $g : E^\nu \rightarrow E^\nu$ such that $\phi = \phi g$, we will prove that g is an automorphism of E^ν . Consider the commutative diagram

$$\begin{array}{ccccc}
 0 & \longrightarrow & M^\nu & \xrightarrow{h} & E \\
 & & \tau_{M^\nu} \cong \downarrow & & \tau_E \cong \downarrow \\
 & & M^{\nu\nu\nu} & \xrightarrow{h^{**}} & E^{\nu\nu} \\
 & & \downarrow \phi^\nu = h^{**\nu} & \nearrow g^* & \\
 & & E^{\nu\nu} & &
 \end{array}$$

We have that $h^{**}\tau_{M^\nu} = g^*\tau_E h$. Therefore, $\tau_E h = g^*\tau_E h$, and $h = \tau_E^{-1}g^*\tau_E h$. Then $\tau_E^{-1}g^*\tau_E$ is an automorphism of E and g^* is an automorphism of $E^{\nu\nu}$. But then g is an automorphism of E^ν . This proves (1) \Rightarrow (2).

In showing (2) \Rightarrow (3) we first show (2) $\Rightarrow R$ is complete. Now let E denote $E(k)$, where $k = R/\mathfrak{m}$. Note that $k \cong \text{Hom}_R(k, E) = k^\nu$ and hence k has a flat cover $F \xrightarrow{\phi} k \rightarrow 0$ by [6, Proposition 1.1]. Since F is flat, F^ν is injective; and both F and F^ν are reflexive. We have the diagram

$$\begin{array}{ccccc}
 0 & \longrightarrow & k^\nu \cong k & \longrightarrow & E \\
 & & \downarrow \phi^* & \nearrow \alpha & \\
 & & F^\nu & &
 \end{array}$$

and α is monic. We can think of E as a submodule of F^ν . Now consider the following commutative diagram with exact rows:

$$\begin{array}{ccccccccc}
 0 & \longrightarrow & E & \xrightarrow{\alpha} & F^\nu & \xrightarrow{\beta} & X & \longrightarrow & 0 \\
 & & \downarrow \tau_E & & \downarrow \tau_{F^\nu} & & \downarrow \tau_X & & \\
 0 & \longrightarrow & E^{\nu\nu} & \xrightarrow{\tilde{\alpha}} & F^{\nu\nu\nu} & \xrightarrow{\tilde{\beta}} & X^{\nu\nu} & \longrightarrow & 0
 \end{array}$$

Since τ_{F^ν} and $\tilde{\beta}$ are surjective, τ_X is surjective and so τ_X is an isomorphism. It follows that τ_E is an isomorphism. Hence E is reflexive. But $R^\nu \cong E$, and since R^ν is reflexive, R is reflexive [4, Proposition 6]. It follows from [9, Theorem 3.7] that R is complete.

Next we show that if R is a complete local ring and every reflexive module has a reflexive flat cover, then $\dim R \leq 1$.

Case 1. Assume that R is a domain. Because E is reflexive, so is F , where

$F \xrightarrow{\phi} E$ is the flat cover of E . We know that ϕ is epic. We have an injection $R \xrightarrow{\cong} E^\nu \xrightarrow{\phi^*} F^\nu$, and F^ν is injective and reflexive. It follows that $E(R)$ is a submodule of F^ν , and so $E(R)$ is reflexive. Let $Q = E(R)$, the field of fraction of R . By Proposition 2.8, $K := Q/R$ is artinian.

If $\mathfrak{m} = 0$ or \mathfrak{m} is nonzero and principal, we are done. So we may assume a_1, a_2, \dots, a_n, b generate \mathfrak{m} , with $n \geq 1$ and $b \neq 0$. Consider, for each i , the descending chain

$$\left(\frac{a_i}{b} + R\right) \supset \left(\frac{a_i^2}{b} + R\right) \supset \cdots \supset \left(\frac{a_i^s}{b} + R\right) \supset \cdots$$

of submodules of K . Since K is an artinian R -module, there exists an integer s such that

$$\left(\frac{a_i^s}{b} + R\right) = \left(\frac{a_i^{s+1}}{b} + R\right).$$

That is,

$$\frac{a_i^s}{b} + R = r \left(\frac{a_i^{s+1}}{b} + R\right)$$

for some $r \in R$, and so

$$\frac{a_i^s}{b} = \frac{ra_i^{s+1}}{b} + c$$

for some $c \in R$. Hence $a_i^s(1 - ra_i) = bc$, and $a_i^s \in (b)$ since $1 - ra_i$ is invertible. From this we can get a sufficiently large integer t so that

$$\mathfrak{m}^t = (Ra_1 + \cdots + Rb)^t \subset (b).$$

So \mathfrak{m} is the radical of a nonzero principal ideal, hence $\dim R = 1$. In [1] this is Theorem 11.14.

Case 2. If R is not a domain, let \mathfrak{p} be a minimal prime ideal of R , and denote the domain (and module) R/\mathfrak{p} by R_1 . Now we are going to prove that the quotient field $Q_1 = E_{R_1}(R_1)$ is a reflexive R_1 -module.

Since R is a reflexive R -module, so is R_1 . If we replace R by R_1 and repeat the argument given in Case 1 mutatis mutandis, then we see that $E_1 := E_R(R_1)$ is a reflexive R -module. But then we know by Lemma 1.1 that

$$Q_1 = \text{ann}_{E_1}(\mathfrak{p}) \subset E_1.$$

It follows that Q_1 is a reflexive R -module, and therefore Q_1 is a reflexive R_1 -module. By Case 1 we know that $\dim R_1 \leq 1$, and so $\dim R \leq 1$.

(3) \Rightarrow (1) By Theorem 2.9.

The last part of the theorem follows by chasing the obvious diagrams, Proposition 3.3, and the fact that for any injective module E , E is a direct summand of $E^{''}$. \square

Acknowledgment

The first author would like to thank the referee for many helpful comments and suggestions. Thanks are also due to Professor E. Enochs for first suggesting this topic, and for many valuable conversations regarding the material in this paper.

References

- [1] M.F. Atiyah and I.G. MacDonald. *Introduction to Commutative Algebra* (Addison-Wesley, Reading, MA, 1969).
- [2] H. Bass. Finitistic dimension and a homological generalization of semi-primary rings, *Trans. Amer. Math. Soc.* 95 (1960) 466–488.
- [3] H. Bass. On the ubiquity of Gorenstein rings, *Math. Z.* 82 (1963) 8–28.
- [4] R. Belshoff. Matlis reflexive modules, *Comm. Algebra* 19 (1991) 1099–1118.
- [5] E. Enochs. Injective and flat covers, envelopes and resolvents, *Israel J. Math.* 39 (1981) 189–209.
- [6] E. Enochs. Flat covers and flat cotorsion modules, *Proc. Amer. Math. Soc.* (2) 92 (1984) 179–184.
- [7] T. Ishikawa. Injective modules and flat modules, *J. Math. Soc. Japan* 17 (1965) 291–296.
- [8] S. Lang. *Algebra* (Addison-Wesley, Menlo Park, CA, 2nd ed., 1984).
- [9] E. Matlis. Injective modules over Noetherian rings, *Pacific J. Math.* 8 (1958) 511–528.
- [10] E. Matlis. Some properties of Noetherian domains of dimension one, *Canad. J. Math.* 13 (1961) 569–586.
- [11] H. Matsumura. *Commutative Ring Theory* (Cambridge University Press, Cambridge, 1989).
- [12] J. Rotman. *An Introduction to Homological Algebra* (Academic Press, New York, 1979).
- [13] D.W. Sharpe and P. Vámos. *Injective Modules* (Cambridge University Press, Cambridge, 1972).